

## **SUPPLEMENTAL MATERIAL**

**Table S1. Outcome at 1 year follow-up, per patient**

Status after 1 year	PTA±BMS (N=64)	DES (N=73)
Alive without amputation or re-intervention	38 (59.4)	47 (64.4)
Major amputation	7 (10.9)	5 (6.8)
Re-intervention	3 (4.7)	3 (4.1)
Amputation and death	4 (6.3)	3 (4.1)
Re-intervention and death	0	1 (1.4)
Re-intervention, amputation and death	1 (1.6)	0
Death	11 (17.2)	14 (19.2)

$P = 0.78 (\chi^2)$ .

BMS indicates bare metal stent; DES, drug-eluting stent; and PTA, percutaneous transluminal angioplasty. Values are N (%).

**Table S2. Outcome at 2 years follow-up, per patient**

Status after 2 years	PTA±BMS (N=64)	DES (N=73)
Alive without amputation or re-intervention	26 (40.6)	42 (57.5)
Major amputation	8 (12.5)	3 (4.1)
Re-intervention	5 (7.8)	4 (5.5)
Amputation and death	5 (7.8)	6 (8.2)
Re-intervention and death	0	1 (1.4)
Re-intervention, amputation and death	1 (1.6)	0
Death	19 (29.7)	17 (23.3)

$P = 0.25 (\chi^2)$

BMS indicates bare metal stent; DES, drug-eluting stent; and PTA, percutaneous transluminal angioplasty. Values are N (%).

**Table S3. Outcome at 3 years follow-up, per patient**

Status after 3 years	PTA±BMS (N=64)	DES (N=73)
Alive without amputation or re-intervention	21 (32.8)	35 (47.9)
Major amputation	8 (12.5)	4 (5.5)
Re-intervention	4 (6.3)	4 (5.5)
Amputation and death	5 (7.8)	6 (8.2)
Re-intervention and death	1 (1.6)	1 (1.4)
Re-intervention, amputation and death	1 (1.6)	0
Death	23 (35.9)	22 (30.1)
Lost to follow-up	1 (1.6)	1 (1.4)

$P = 0.61 (\chi^2)$

BMS indicates bare metal stent; DES, drug-eluting stent; and PTA, percutaneous transluminal angioplasty. Values are N (%).

**Table S4. Outcome at 4 years follow-up, per patient**

Status after 4 years	PTA±BMS (N=64)	DES (N=73)
Alive without amputation or re-intervention	12 (18.8)	29 (39.7)
Major amputation	7 (10.9)	1 (1.4)
Re-intervention	2 (3.1)	4 (5.5)
Amputation and death	5 (7.8)	7 (9.6)
Re-intervention and death	1 (1.6)	1 (1.4)
Re-intervention, amputation and death	1 (1.6)	0
Death	29 (45.3)	24 (32.9)
Lost to follow-up	1 (1.6)	2 (2.7)
Follow-up ongoing	4 (6.3)	3 (4.1)
Amputation, follow-up ongoing	2 (3.1)	2 (2.7)

$P=0.12 (\chi^2)$

BMS indicates bare metal stent; DES, drug-eluting stent; and PTA, percutaneous transluminal angioplasty. Values are N (%).

**Table S5. Outcome at 5 years follow-up, per patient**

Status after 5 years	PTA±BMS (N=64)	DES (N=73)
Alive without amputation or re-intervention	8 (12.5)	11 (15.1)
Major amputation	5 (7.8)	2 (2.7)
Re-intervention	1 (1.6)	3 (4.1)
Amputation and death	5 (7.8)	7 (9.6)
Re-intervention and death	1 (1.6)	2 (2.7)
Re-intervention, amputation and death	1 (1.6)	0
Death	31 (48.4)	32 (43.8)
Lost to follow-up	1 (1.6)	4 (5.5)
Follow-up ongoing	5 (7.8)	9 (12.3)
Re-intervention, follow-up ongoing	1 (1.6)	1 (1.4)
Amputation, follow-up ongoing	5 (7.8)	2 (2.7)

$P=0.63$  ( $\chi^2$ )

BMS indicates bare metal stent; DES, drug-eluting stent; and PTA, percutaneous transluminal angioplasty. Values are N (%).