

Patients With Heart Failure Readmitted to the Original Hospital Have Better Outcomes Than Those Readmitted Elsewhere

Finlay A. McAlister, MD, MSc; Erik Youngson, MMath; Padma Kaul, PhD

Background—Up to one fifth of readmissions after a heart failure hospitalization occur at a different hospital. This negatively impacts information continuity, but whether site of readmission impacts subsequent outcomes is unclear.

Methods and Results—Retrospective cohort study of all patients discharged with a primary diagnosis of heart failure in Canada between April 2004 and December 2013. We compared patients readmitted within 30 days to the original hospital versus a different hospital. Of the 217 039 heart failure patients (mean age, 76.8 years, 50.1% male), 39 368 (18.1%) were readmitted within 30 days—32 771 (83.2%) to the original hospital and 6597 (16.8%) to a different hospital (increasing over time from 15.6% in 2004 to 18.5% by 2013; P for trend=0.001). Patients readmitted to different hospitals were younger and were more likely to be male, have a rural residence, a more-recent discharge year, an index hospitalization at a teaching hospital, and to be brought in by ambulance at the time of the readmission. Readmissions to the original hospital were substantially shorter (mean, 10.4 days [95% CI, 10.3–10.6] versus 11.6 days [95% CI, 11.3–12.0]; adjusted means, 11.0 versus 12.0; P <0.0001) and had lower mortality (14.4% versus 15.0%; adjusted odds ratio, 0.89; 95% CI, 0.82–0.96) than readmissions to different hospitals.

Conclusions—Readmissions to a different hospital are becoming more frequent over time and are associated with longer stays and higher mortality rates than readmissions to the original hospital. Our findings provide further evidence that care fragmentation may be deleterious for patients with heart failure. (*J Am Heart Assoc.* 2017;6:e004892. DOI: 10.1161/JAHA.116.004892.)

Key Words: heart failure • outcomes research • readmissions

Heart Failure (HF) is one of the most common reasons for hospitalization in North America,¹ and the risks of hospitalization and readmission are continuing to increase.^{2,3} Although most patients are readmitted to the hospital they were originally discharged from, approximately one fifth are readmitted to a different hospital.^{4–8} Because most hospitals do not share inpatient data (such as admission and progress notes, diagnostic test results, and in-hospital therapies or interventions) with one another⁹ and given the well-known deficiencies in discharge summary completeness and promptness,¹⁰ readmission to a different hospital negatively impacts information continuity.

Whether site of readmission impacts subsequent outcomes is unclear. Whereas 1 study⁴ suggested that 30-day readmissions to a different hospital were associated with higher mortality, another⁵ did not find any difference in

mortality rates for those readmitted to the original versus a different hospital. However, both studies evaluated patients discharged after hospitalization for any reason—large proportions of which were surgical hospitalizations—and it is already known that surgical patients demonstrate markedly poorer outcomes when readmitted to a different hospital.⁷ Thus, the influence of readmission site for patients with HF as the primary diagnosis for their index hospitalization is unknown. Although this is an important policy question with implications for clinicians, patients, ambulance referral practices, and hospital funding models, it has not been addressed in a randomized trial. Thus, we designed a population-based cohort study to test whether site of hospital readmission influenced subsequent length of stay or in-patient mortality for patients with a primary diagnosis of HF.

From the Division of General Internal Medicine (F.A.M.) and Patient Health Outcomes Research and Clinical Effectiveness Unit (F.A.M., E.Y.), University of Alberta, Edmonton, Alberta, Canada; Canadian VIGOUR Centre (F.A.M., P.K.) and Division of Cardiology (P.K.), Mazankowski Alberta Heart Institute, Edmonton, Alberta, Canada.

Correspondence to: Finlay A. McAlister, MD, MSc, 5-134C Clinical Sciences Building, University of Alberta, 11350 83 Ave, Edmonton, Alberta, Canada T6G 2G3. E-mail: finlay.mcalister@ualberta.ca

Received October 18, 2016; accepted March 21, 2017.

© 2017 The Authors. Published on behalf of the American Heart Association, Inc., by Wiley. This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

Methods

Setting

Canada has a system of universal health insurance that provides access to medical care at no direct cost to the patient, and all acute care hospitals in Canada submit information to the Discharge Abstract Database held by the Canadian Institute for Health Information.

Study Sample

We used the Discharge Abstract Database in 9 Canadian provinces and 3 territories to examine all acute care hospitalizations with a primary diagnosis of HF using International Classification of Diseases, Tenth Revision code I50.x (93% positive predictive value for HF when validated against chart audit)¹¹ between April 1, 2004 and December 31, 2013. The Discharge Abstract Database includes the admission and discharge dates, acuity (elective versus urgent/emergent), the primary diagnosis assigned by the attending physician, and up to 24 other diagnoses coded by trained nosologists in each hospital using nationally standardized definitions.

Hospitalizations that overlapped within 24 hours were combined into 1 episode of care (which was the unit of analysis), and the index hospital was defined as the last hospital in the episode of care cascade that had HF as the primary diagnosis. Length of stay (LOS) was summed across all hospitals for each episode of care. For patients with multiple HF episodes during the study time period, we selected their first hospitalization as the index.

In the primary analysis, we analyzed a cohort of adult patients discharged alive after a typical hospitalization for HF; thus, we excluded patients who were younger than 20 years old, died during the index hospitalization, were hospitalized for longer than 30 days, or signed themselves out against medical advice. We identified any nonelective readmissions for any cause between 1 and 30 days after discharge (as mentioned previously, we excluded readmissions within 24 hours of discharge to avoid misattributing transfers between hospitals and instead combined these with the initial hospitalization into a single episode of care). For patients with multiple readmissions within 30 days of the index hospitalization, we used the first readmission for analysis.

Covariates

Comorbidities for each patient were identified using International Classification of Diseases, Tenth Revision codes from the Discharge Abstract Database for the index hospitalization and any hospitalizations in the previous 12 months.¹² Other covariates included type of hospital (teaching, large community, medium community, and small community), patient

residence (rural versus urban), number of hospitalizations in the past 6 months, day of discharge (given that this is known to influence readmission rates),¹³ and specialty of most responsible physician during the index hospitalization (cardiologist, other internal medicine, family physician, or other).

Outcomes

Our primary outcomes of interest were LOS and in-hospital mortality for HF patients readmitted for any cause within 30 days to the original versus a different hospital. In a sensitivity analysis, we examined these outcomes in only those patients readmitted for HF to the original versus a different hospital.

Statistical Analyses

Ages and patient comorbidities were compared using *t* tests or chi-squared tests between those patients who were/were not readmitted within 30 days and between those who were readmitted to the original versus a different hospital. We identified which factors were associated with different versus original hospital readmission using multivariable logistic regression adjusting for age, sex, Charlson score, number of hospitalizations in past 6 months, acuity of admission, type of most responsible physician during index hospitalization, discharge disposition after index hospitalization, day of discharge from index hospitalization, type of hospital for index hospitalization, rural residence for the patient, day of readmission (weekday versus weekend/holiday), arrival by ambulance to readmission hospital, year of index hospitalization, and 2 random intercepts to account for clustering effects of province and of hospital. A similar logistic regression model including the same covariates was used to obtain an adjusted odds ratio comparing in-hospital mortality between patients who were readmitted to the original versus a different hospital. To compare the readmission LOS between patients readmitted to the original versus a different hospital, a generalized linear mixed model was used to obtain adjusted length of stay by least squares means and adjusting for age, sex, rural residence, Charlson score, type of hospital, type of most responsible physician during index hospitalization, discharge disposition after index hospitalization, day of readmission, arrival by ambulance to readmission hospital, and 2 random intercepts (thus the mixed model) to account for clustering effects of province and hospital. A model using log-transformed LOS was also examined, but did not appreciably alter the results. SAS software (version 9.4; SAS Institute Inc, Cary, NC) was used for all statistical analysis. The University of Alberta Health Research Ethics Board approved this study with waiver of informed consent because we were using de-identified data.

Results

Of the 217 039 patients (mean age, 76.8 years; 50.1% male) discharged alive after an acute care hospitalization for HF (Figure), 39 368 (18.1%) were readmitted within 30 days—32 771 (83.2%) to the original hospital and 6597 (16.8%) to a different hospital. There was a statistically significant increase in the proportion being readmitted to a different hospital over time: from 15.6% in 2004 to 18.5% in 2013 (*P* for trend=0.001). Patients were more likely to be readmitted to the original hospital if they were older, female, lived in an urban setting, had shorter LOS during the index hospitalization, had lower comorbidity burdens, were treated in a community hospital, and if their attending physician during the index hospitalization had been a primary care physician (Table 1)—however, many of these differences disappeared on multivariate analysis. The factors independently associated with being readmitted to a different hospital (after adjusting for Charlson scores and the other factors in Table 1) were male sex, younger age, rural residence, more-recent discharge year, long-term care facility resident, index hospitalization was at a teaching hospital, and being brought in by ambulance for the readmission; patients cared for by family physicians were less likely to be readmitted to a different hospital (Table 2).

The top 5 causes for readmission in this cohort of patients were HF (n=14 535; 36.9%), chronic obstructive pulmonary

disease (n=1945; 4.9%), acute myocardial infarction (n=1332; 3.4%), pneumonia (n=1133; 2.9%), and atrial fibrillation or flutter (n=940; 2.4%).

Whereas readmitted patients exhibited longer LOS during the readmission than during their index hospitalization (mean, 8.2 days; 95% CI, 8.1–8.2), those readmitted to the original hospital demonstrated substantially shorter LOS for the readmission (mean, 10.4 days; 95% CI, 10.3–10.6; adjusted mean, 11.0; 95% CI, 10.1–12.0) than those readmitted to different hospitals (mean, 11.6 days; 95% CI, 11.3–12.0; adjusted mean, 12.0; 95% CI, 11.1–13.0; *P*<0.0001). Patients readmitted to the original hospital also demonstrated lower mortality rates during the readmission (14.4% versus 15.0% for those readmitted to a different hospital; adjusted odds ratio, 0.89; 95% CI, 0.82–0.96). Of note, perusal of our exclusion criteria (Figure) reveals that mortality during the index hospitalization was 6.1%.

Our sensitivity analysis restricted to patients readmitted for HF as the primary diagnosis found similar patterns in terms of factors associated with readmission to the original hospital versus a different hospital and worse outcomes for readmissions than index hospitalizations.

Discussion

We found that readmission to a different hospital was common for patients with HF, was becoming more frequent over time, and was associated with longer LOS and higher inpatient mortality rates than readmissions to the original hospital. Although we could not examine resource use in our data set, a recent Medicare study found that patient readmissions to different hospitals generated higher costs because of more diagnostic testing and procedures than readmissions to the original hospital.⁵ Although it is already known¹ that readmissions are associated with longer LOS and poorer outcomes than the original hospitalization, our study adds to the literature base by demonstrating that readmissions to different hospitals have even worse outcomes than readmissions to the original hospital. Our study also helps clarify the conflicting results from 2 previous studies of this phenomenon: ¹⁴ demonstrated higher mortality rates with different hospital readmissions whereas the other⁵ did not. Furthermore, as mentioned in the Introduction, both previous studies evaluated a broad mix of patients discharged after hospitalization for a wide variety of medical or surgical diagnoses, and neither examined LOS during the readmission.

We cannot say definitively why alternate hospital readmissions have poorer outcomes because this is an observational study. However, we theorize that fragmentation of care plays a role. Previous studies^{14,15} have shown better outcomes for




Figure. Cohort derivation. *Transferred to inpatient facility as final discharge disposition, but no subsequent hospitalization within 24 hours. HF indicates heart failure; LOS, length of stay.

Table 1. Patient Characteristics at the Time of the Index Hospitalization With a Primary Diagnosis of HF

Characteristic	Not Readmitted Within 30 Days n=177 671	Readmitted Within 30 Days n=39 368	P Value	Readmitted Within 30 Days to Original Hospital n=32 771	Readmitted Within 30 Days to Different Hospital n=6597	P Value
Age at time of index hospitalization	76.6 (12.3)	77.6 (11.5)	<0.0001	78.1 (11.3)	75.1 (12.5)	<0.0001
Male	88 770 (50.0)	19 954 (50.7)	0.10	16 361 (49.9)	3593 (54.5)	<0.0001
Rural resident	37 362 (21.1)	8680 (22.1)	<0.0001	6886 (21.0)	1794 (27.2)	<0.0001
Admitted on a nonholiday weekday*	130 084 (73.2)	28 450 (72.3)	0.0001	23 630 (72.1)	4764 (72.2)	0.86
Arrived by ambulance*	75 798 (42.7)	18 079 (45.9)	<0.0001	17 338 (52.9)	3595 (54.5)	0.02
Resided in long-term care facility before index hospitalization	15 672 (8.8)	3583 (9.1)	0.08	3136 (9.6)	447 (6.8)	<0.0001
Charlson comorbidity index score	4.0 (2.1)	4.5 (2.3)	<0.0001	4.5 (2.3)	4.6 (2.3)	<0.0001
Hospitalizations during previous 6 months	0.3 (0.7)	0.5 (0.9)	<0.0001	0.5 (0.9)	0.5 (0.9)	0.10
Acute length of stay for index hospitalization	7.7 (5.9)	8.2 (6.2)	<0.0001	8.0 (6.0)	8.9 (6.8)	<0.0001
LACE score	11.4 (2.5)	11.9 (2.5)	<0.0001	11.8 (2.5)	12.0 (2.6)	<0.0001
Discharged on weekend/holiday	27 633 (15.6)	6247 (15.9)	0.12	5229 (16.0)	1018 (15.4)	0.29
Discharge disposition			<0.0001			<0.0001
Home	121 425 (68.3)	25 587 (65.0)		21 116 (64.4)	4471 (67.8)	
Home with homecare	34 046 (19.2)	9775 (24.8)		8352 (25.5)	1423 (21.6)	
Long-term care/skilled nursing facility	22 200 (12.5)	4006 (10.2)		3303 (10.1)	703 (10.7)	
Past history of heart failure in the 1 year before index	28 744 (16.2)	9333 (23.7)	<0.0001	7853 (24.0)	1480 (22.4)	0.008
Hypertension	73 199 (41.2)	16 705 (42.4)	<0.0001	13 581 (41.4)	3124 (47.4)	<0.0001
Diabetes mellitus	63 276 (35.6)	15 319 (38.9)	<0.0001	12 692 (38.7)	2627 (39.8)	0.10
Ishemic heart disease, including past MI, CABG, or PCI/stent	64 634 (36.4)	16 191 (41.1)	<0.0001	13 223 (40.3)	2968 (45.0)	<0.0001
Peripheral arterial disease	1203 (0.7)	357 (0.9)	<0.0001	284 (0.9)	73 (1.1)	0.06
Cerebrovascular disease (previous stroke/TIA)	3265 (1.8)	854 (2.2)	<0.0001	689 (2.1)	165 (2.5)	0.04
Atrial fibrillation/flutter	61 682 (34.7)	14 111 (35.8)	<0.0001	11 719 (35.8)	2392 (36.3)	0.44
Ventricular arrhythmias	3593 (2.0)	916 (2.3)	0.0001	726 (2.2)	190 (2.9)	0.0011
Chronic kidney disease	37 805 (21.3)	11 224 (28.5)	<0.0001	9176 (28.0)	2048 (31.0)	<0.0001
Cancer	7028 (4.0)	2267 (5.8)	<0.0001	1858 (5.7)	409 (6.2)	0.09
COPD or asthma	35 721 (20.1)	9421 (23.9)	<0.0001	7902 (24.1)	1519 (23.0)	0.06
Peptic ulcer disease	1650 (0.9)	480 (1.2)	<0.0001	376 (1.1)	104 (1.6)	0.004
Liver disease	2145 (1.2)	639 (1.6)	<0.0001	516 (1.6)	123 (1.9)	0.09
Dementia	9621 (5.4)	2106 (5.3)	0.60	1764 (5.4)	342 (5.2)	0.51
Index hospital characteristics			<0.0001			<0.0001
Teaching hospital	41 136 (23.3)	8815 (22.5)		6675 (20.5)	2140 (32.7)	
Large community hospital	78 424 (44.4)	16 837 (43.0)		14 278 (43.8)	2559 (39.0)	
Medium community hospital	34 604 (19.6)	7671 (19.6)		6730 (20.6)	941 (14.4)	
Small community hospital	22 618 (12.8)	5824 (14.9)		4910 (15.1)	914 (13.9)	
Most responsible physician during index hospitalization			<0.0001			<0.0001
Cardiologist	30 880 (17.4)	5717 (14.5)		4252 (13.0)	1465 (22.2)	

Continued

Table 1. Continued

Characteristic	Not Readmitted Within 30 Days n=177 671	Readmitted Within 30 Days n=39 368	P Value	Readmitted Within 30 Days to Original Hospital n=32 771	Readmitted Within 30 Days to Different Hospital n=6597	P Value
Internal medicine other than cardiology	50 576 (28.5)	11 404 (29.0)		9119 (27.8)	2285 (34.6)	
Family physician	85 386 (48.1)	19 778 (50.2)		17 393 (53.1)	2385 (36.2)	
Other	10 829 (6.1)	2469 (6.3)		2007 (6.1)	462 (7.0)	

Data are presented as means (SD) or n (%). P values calculated using t test (means) or chi-square test (binary/categorical). Missing values: 337 (0.16%) for rural residence status; 1110 (0.51%) for hospital type. LACE score=index to identify risk for readmission or death, L stands for length of stay, A stands for acuity of admission, C stands for Charlson comorbidity score, and E stands for number of emergency department visits in the past 6 months. CABG indicates coronary artery bypass grafting; COPD, chronic obstructive pulmonary disease; MI, myocardial infarction; PCI, percutaneous coronary intervention; TIA, transient ischemic attack.

*Refers to the index hospitalization for the first 2 columns and the readmission hospitalization for the last 2 columns.

HF patients cared for by physicians familiar with them after hospital discharges or emergency department visits. Other studies have also demonstrated that continuity of physician care reduces hospitalizations, especially for patients with high comorbidity burdens.¹⁶ Thus, it seems reasonable to hypothesize that continuity of inpatient care would be associated with improved outcomes—however, we acknowledge that this is a hypothesis that requires testing in a prospective, controlled study.

Although it is possible that our results are a manifestation of residual confounding, it is worth noting that patients readmitted to a different hospital were not only younger and less likely to have been urgently admitted during the index hospitalization (suggesting that they were perhaps less sick), but they were also more likely to have been treated by a specialist or in an academic teaching hospital during their preceding index event—all factors shown to be associated with better subsequent outcomes.¹³ This would suggest that any residual confounding may, in fact, have biased toward the null in our study by improving outcomes in the “different hospital readmission” group. Regardless, without information on the factors motivating alternate hospital readmissions, we cannot comment on the potential modifiability of this practice. However, given the poorer prognosis associated with alternate hospital readmission, our data suggest that clinicians should counsel patients on the potential benefits of returning to their original hospital if they deteriorate. In the same vein, we believe that our findings should prompt policy makers to revisit current ambulance policies which focus on shortening wait times in emergency rooms by diverting ambulances to the least busy emergency room regardless of where they had been most recently discharged from. In fact, given that delaying care so that one can be cared for by familiar providers may have harms as well as benefits, we believe a controlled trial comparing sending patients back to the original hospital versus to the closest facility should be conducted. Our findings also have implications for hospital performance measures and funding models based on

readmission rates given the substantial proportion of patients readmitted to hospitals other than the one they were discharged from. Finally, our results should motivate attempts to improve the sharing of patient information between hospitals—one of the many hopes for electronic medical records.

Despite reporting outcomes in all adults (ie, not just elderly patients) hospitalized for HF in 9 provinces and 3 territories of Canada (a country with free universal access to acute care hospitals), robust data permitting adjustment for comorbidities as well as health resource use, and a large sample, there are some limitations to our analysis. First, we relied on administrative data to define HF and comorbidities and do not have information on ejection fraction, natriuretic peptide levels, nor clinical findings that would permit delineation of illness severity. Although we acknowledge this weakness, we used validated International Classification of Diseases codes and data definition algorithms^{11,12} to build comorbidity profiles and the outcomes we evaluated (all-cause rehospitalizations, LOS, and in-hospital mortality) are relevant in HF patients regardless of ejection fraction, etiology, or clinical status. Second, although outpatient visits, particularly with familiar physicians^{14–16} or access to specialized clinics¹⁷ after discharge, would reduce readmissions, we do not have any data on whether availability of these outpatient resources varied over time or whether outpatient physician visits were or were not with physicians associated with the original hospital. Third, we had no data on distance from patient home to nearest hospital nor socioeconomic factors, such as homelessness or access to motor vehicle, that may have influenced both health outcomes and choice of hospital. Fourth, we had no data on completeness, promptness, or dissemination of discharge summaries after the index hospitalization, although previous studies suggest that these are low within a 30-day time frame.¹⁸ Finally, although we focused on 30-day readmissions, a study in 1 Canadian province suggested that mortality differences observed at 30 days persisted across a full year of follow-up.⁴

Table 2. Factors Associated With Readmission to a Hospital Different Than the Index Hospital

Variable	Univariate Odds Ratio (95% CI)	Adjusted Odds Ratio (95% CI)
Age (per 1 year increase)	0.98 (0.98, 0.98)	0.98 (0.98, 0.98)
Male	1.20 (1.14, 1.27)	1.10 (1.04, 1.16)
Rural residence	2.42 (2.23, 2.63)	2.43 (2.23, 2.65)
Charlson Score (per 1-unit increase)	1.01 (1.00, 1.03)	1.00 (0.99, 1.01)
Number of hospitalizations in previous 6 months (per 1-unit increase)	1.03 (1.00, 1.07)	1.02 (0.99, 1.05)
Urgent index admission (vs elective)	0.55 (0.45, 0.68)	0.61 (0.50, 0.75)
Family physician as most responsible during index	0.55 (0.50, 0.59)	0.61 (0.55, 0.66)
Index discharge on weekend/holiday	0.97 (0.90, 1.04)	0.94 (0.87, 1.02)
Year of index discharge (per 1-year increase)	1.01 (1.00, 1.02)	1.02 (1.01, 1.03)
Discharge disposition		
Home (ref)	1 (ref)	1 (ref)
Home with homecare	0.77 (0.71, 0.82)	0.89 (0.82, 0.95)
Long term care/skilled nursing facility	0.95 (0.87, 1.05)	1.17 (1.06, 1.28)
Index hospital type		
Teaching (ref)	1 (ref)	1 (ref)
Large community	0.63 (0.52, 0.77)	0.74 (0.60, 0.92)
Medium community	0.52 (0.42, 0.65)	0.58 (0.46, 0.74)
Small community	0.66 (0.54, 0.81)	0.59 (0.46, 0.75)
Readmission on weekend/holiday	1.02 (0.96, 1.08)	1.02 (0.96, 1.09)
Arrived by ambulance to readmission hospital	1.07 (1.01, 1.13)	1.22 (1.15, 1.29)

Adjusted odds ratios obtained from model including all variables listed in table. Response variable=readmission to NON-Index hospital (vs readmission to same hospital as index admission). Eg, odds ratio >1 implies positive association with readmission to different hospital (compared to readmission to same hospital).

In conclusion, our finding that readmissions to hospitals other than the original one are associated with poorer outcomes for patients with HF reinforces concerns that such readmissions may not only result in duplication of tests and procedures⁵ and lead to exposure to potentially different nosocomial pathogens,⁴ but also may delay timely diagnoses or treatments. We believe our study provides further support for the importance of continuity in health care and should motivate patients, caregivers, and their physicians to ensure follow-up after discharge occurs with healthcare providers who

are familiar with them. Our study should also motivate health system planners to conduct evaluations of “return to original hospital” versus “take to the closest facility” ambulance policies for HF patients who deteriorate post-discharge.

Sources of Funding

Dr McAlister receives salary support from Alberta Innovates Health Solution (AIHS) and holds the University of Alberta Chair in Cardiovascular Outcomes Research.

Disclosures

All authors declare that they have no financial interests that may be relevant to the submitted work. Dr McAlister is guarantor of this work.

References

- Jencks SF, Williams MV, Coleman EA. Rehospitalizations among patients in the Medicare fee-for-service program. *N Engl J Med*. 2009;360:1418–1428.
- Go AS, Mozaffarian D, Roger VL, Benjamin EJ, Berry JD, Borden WB, Bravata DM, Dai S, Ford ES, Fox CS, Franco S, Fullerton HJ, Gillespie C, Hailpern SM, Heit JA, Howard VJ, Huffman MD, Kissela BM, Kittner SJ, Lackland DT, Lichtman JH, Lisabeth LD, Magid D, Marcus GM, Marelli A, Matchar DB, McGuire DK, Mohler ER, Moy CS, Mussolino ME, Nichol G, Paynter NP, Schreiner PJ, Sorlie PD, Stein J, Turan TN, Virani SS, Wong ND, Woo D, Turner MB; American Heart Association Statistics Committee and Stroke Statistics Subcommittee. Executive summary: heart disease and stroke statistics—2013 update: a report from the American Heart Association. *Circulation*. 2013;127:143–152.
- Bueno H, Ross JS, Wang Y, Chen J, Vidán MT, Normand SL, Curtis JP, Drye EE, Lichtman JH, Keenan PS, Kosiborod M, Krumholz HM. Trends in length of stay and short-term outcomes among Medicare patients hospitalized for heart failure, 1993–2006. *JAMA*. 2010;303:2141–2247.
- Staples JA, Thiruchelvam D, Redelmeier DA. Site of hospital readmission and mortality: a population-based retrospective cohort study. *CMAJ Open*. 2014;2:E77–E85.
- Kind AJM, Bartels C, Mell MW, Mullahy J, Smith M. For-profit hospital status and rehospitalizations at different hospitals: an analysis of Medicare data. *Ann Intern Med*. 2010;153:718–727.
- Hempstead K, DeLia D, Cantor JC, Nguyen T, Brenner J. The fragmentation of hospital use among a cohort of high utilizers. Implications for emerging care coordination strategies for patients with multiple chronic conditions. *Med Care*. 2014;52:S67–S74.
- Tsai TC, Orav EJ, Jha AK. Care fragmentation in the postdischarge period. Surgical readmissions, distance of travel, and postoperative mortality. *JAMA Surg*. 2015;150:59–64.
- Bourgeois FC, Olson KL, Mandl KD. Patients treated at multiple acute health care facilities. Quantifying information fragmentation. *Arch Intern Med*. 2010;170:1989–1995.
- Adler-Milstein J, DesRoches CM, Jha AK. Health information exchange among US hospitals. *Am J Manag Care*. 2011;17:761–768.
- Kripalani S, LeFevre F, Phillips CO, Williams MV, Basaviah P, Baker DW. Deficits in communication and information transfer between hospital-based and primary care physicians. *JAMA*. 2007;297:831–841.
- Quan H, Li B, Saunders LD, Parsons GA, Nilsson CI, Alibhai A, Ghali WA; for the IMECCHI Investigators. Assessing validity of ICD-9-CM and ICD-10 administrative data in recording clinical conditions in a unique dually coded database. *Health Serv Res*. 2008;43:1424–1441.
- Lee DS, Donovan L, Austin PC, Gong Y, Liu PP, Rouleau JL, Tu JV. Comparison of coding of heart failure and comorbidities in administrative and clinical data for use in outcomes research. *Med Care*. 2005;43:182–188.
- McAlister FA, Au A, Majumdar SR, Youngson E, Padwal RS. Postdischarge outcomes in heart failure are better for teaching hospitals and weekday discharges. *Circ Heart Fail*. 2013;6:922–929.

14. McAlister FA, Youngson E, Bakal JA, Kaul P, Ezekowitz J, van Walraven C. Impact of physician continuity on death or urgent readmission after discharge among patients with heart failure. *CMAJ*. 2013;185:e681–e689.
15. Sidhu R, Youngson E, McAlister FA. Physician continuity improves outcomes for heart failure patients treated and released from the emergency department. *JACC Heart Fail*. 2014;2:368–376.
16. Gruneir A, Bronskill SE, Maxwell CJ, Bai YQ, Kone AJ, Thavorn K, Petrosyan Y, Calzavara A, Wodchis WP. The association between multimorbidity and hospitalization is modified by individual demographics and physician continuity of care: a retrospective cohort study. *BMC Health Serv Res*. 2016;16:154.
17. McAlister FA, Stewart S, Ferrua S, McMurray JJV. Multidisciplinary strategies for the management of heart failure patients at high risk for admission: a systematic review of randomized trials. *J Am Coll Cardiol*. 2004;44:810–819.
18. van Walraven C, Taljaard M, Bell CM, Etchells E, Stiell IG, Zarnke K, Forster AJ. A prospective cohort study found that provider and information continuity was low after patient discharge from hospital. *J Clin Epidemiol*. 2010;63:1000–1010.



Patients With Heart Failure Readmitted to the Original Hospital Have Better Outcomes Than Those Readmitted Elsewhere

Finlay A. McAlister, Erik Youngson and Padma Kaul

J Am Heart Assoc. 2017;6:e004892; originally published May 10, 2017;
doi: 10.1161/JAHA.116.004892

The *Journal of the American Heart Association* is published by the American Heart Association, 7272 Greenville Avenue, Dallas, TX 75231
Online ISSN: 2047-9980

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://jaha.ahajournals.org/content/6/5/e004892>